## CS 70 Discrete Mathematics and Probability Theory Dis 05B

## 1 Head Count

Consider a coin with  $\mathbb{P}(\text{Heads}) = 2/5$ . Suppose you flip the coin 20 times, and define *X* to be the number of heads.

(a) Name the distribution of *X* and what its parameters are.

(b) What is  $\mathbb{P}(X = 7)$ ?

(c) What is  $\mathbb{P}(X \ge 1)$ ? Hint: You should be able to do this without a summation.

(d) What is  $\mathbb{P}(12 \le X \le 14)$ ?

## 2 How Many Queens?

You shuffle a standard 52-card deck, before drawing the first three cards from the top of the pile. Let X denote the number of queens you draw.

(a) What is  $\mathbb{P}(X=0)$ ,  $\mathbb{P}(X=1)$ ,  $\mathbb{P}(X=2)$  and  $\mathbb{P}(X=3)$ ?

(b) What do your answers you computed in part a add up to?

(c) Compute  $\mathbb{E}(X)$  from the definition of expectation.

(d) Let  $X_i$  be an indicator random variable that equals 1 if the *i*th card a is queen and 0 otherwise. Are the  $X_i$  indicators independent?

## 3 Linearity

Solve each of the following problems using linearity of expectation. Explain your methods clearly.

(a) In an arcade, you play game A 10 times and game B 20 times. Each time you play game A, you win with probability 1/3 (independently of the other times), and if you win you get 3 tickets (redeemable for prizes), and if you lose you get 0 tickets. Game B is similar, but you win with probability 1/5, and if you win you get 4 tickets. What is the expected total number of tickets you receive?

(b) A monkey types at a 26-letter keyboard with one key corresponding to each of the lower-case English letters. Each keystroke is chosen independently and uniformly at random from the 26 possibilities. If the monkey types 1 million letters, what is the expected number of times the sequence "book" appears?